

# TABLE OF CONTENTS

---

|                  |  |               |
|------------------|--|---------------|
| <b>Section 5</b> | <b>Environmental Information</b> .....             | <b>5.16-1</b> |
| 5.16             | Paleontological Resources .....                    | 5.16-1        |
| 5.16.1           | Affected Environment.....                          | 5.16-3        |
| 5.16.1.1         | Geographic Location .....                          | 5.16-3        |
| 5.16.1.2         | Regional Geologic Setting.....                     | 5.16-4        |
| 5.16.1.3         | Resource Inventory Methods.....                    | 5.16-5        |
| 5.16.1.4         | Paleontological Resource Assessment Criteria ..... | 5.16-5        |
| 5.16.1.5         | Resource Inventory Results .....                   | 5.16-10       |
| 5.16.2           | Environmental Consequences.....                    | 5.16-13       |
| 5.16.3           | Cumulative Impacts Analyses.....                   | 5.16-14       |
| 5.16.4           | Mitigation Measures .....                          | 5.16-14       |
| 5.16.5           | Laws, Ordinances, Regulations, and Standards ..... | 5.16-15       |
| 5.16.5.1         | Federal .....                                      | 5.16-16       |
| 5.16.5.2         | State .....  | 5.16-16       |
| 5.16.5.3         | Local .....  | 5.16-16       |
| 5.16.6           | Involved Agencies and Agency Contacts .....        | 5.16-17       |
| 5.16.7           | Permits Required and Permit Schedule.....          | 5.16-17       |
| 5.16.8           | References.....                                    | 5.16-17       |

## Tables

|              |   |
|--------------|---|
| Table 5.16-1 | Summary of LORS – Paleontological Resources |
|--------------|---|

# TABLE OF CONTENTS

---

*This page intentionally left blank*

## 5.16 PALEONTOLOGICAL RESOURCES

Hydrogen Energy International LLC (HEI or Applicant) is jointly owned by BP Alternative Energy North America Inc., and Rio Tinto Hydrogen Energy LLC. HEI is proposing to build an Integrated Gasification Combined Cycle (IGCC) power generating facility called Hydrogen Energy California (HECA or the “Project”) in Kern County, California. The Project will produce electricity while substantially reducing greenhouse gas emissions by capturing carbon dioxide (CO<sub>2</sub>) and transporting it for enhanced oil recovery (EOR) and sequestration.

The 315-acre Project Site is located approximately 6.5 miles west of the outermost edge of the city of Bakersfield and 2 miles northwest of the unincorporated community of Tupman in western Kern County, California, as shown in Figure 2-1, Project Vicinity Map. The Project Site is adjacent to an oil producing area known as the Elk Hills Oil Field Unit. The Project Site is currently undeveloped. Existing surface elevations vary from about 445 feet above mean sea level (msl) in the southwest corner to about 310 feet above msl in the northeast corner.

The Project will gasify petroleum coke (or blends of petroleum coke and coal, as needed) to produce hydrogen to fuel a combustion turbine operating in combined cycle mode. The gasification component feeds a 390 gross megawatt (MW) combined cycle plant. The net electrical generation output from the Project will provide California with approximately 250 MW of low-carbon baseload power to the grid. The gasification component will also capture approximately 90 percent of the carbon dioxide from the syngas at steady-state operation, which will be transported and used for EOR and sequestration (storage) in the Elk Hills Oil Field Unit. In addition, approximately 100 MW of natural gas generated peaking power will be available from the Project.

The Project Site and linear facilities comprise the affected study area and are entirely located in Kern County, California. These Project components are described below.

Major on-site Project components will include, as shown on Figure 2-4, Plot Plan:

- Solids Handling, Gasification, and Gas Treatment
  - Feedstock delivery, handling and storage
  - Gasification
  - Sour shift/gas cooling
  - Mercury removal
  - Acid gas removal
- Power Generation
  - Combined-cycle power generation
  - Auxiliary combustion turbine generator
  - Electrical switching facilities
- Supporting Process Systems
  - Natural gas fuel systems

- Air separation unit (ASU)
- Sulfur recovery unit
- Zero liquid discharge
- Carbon dioxide compression
- Wastewater injection wells
- Raw water treatment plant
- Other plant systems

The Project also includes the following off-site facilities, as shown on Figure 2-5, Project Location Map:

- **Electrical Transmission Line** – An electrical transmission line will interconnect the Project to Pacific Gas & Electric’s (PG&E) Midway Substation. The interconnection voltage is expected to be 230 kilovolts (kV). The Project is considering two alternative transmission routes, both of which extend from the western edge of the Project Site to the north, and west to the north side of the substation. Transmission Alternative 1 is approximately 9 miles long and Transmission Alternative 2 is approximately 9.5 miles long.
- **Natural Gas Supply** – A natural gas interconnection will be made with either PG&E or Southern California Gas Company natural gas pipelines, both of which are located southeast of the Project Site. The natural gas pipeline will be approximately 7 miles in length. The interconnect will consist of one tap off the existing natural gas line, one meter set, one service pipeline service connection, and a pressure limiting station located on the Project Site.
- **Water Supply Pipelines** – The Project will utilize brackish groundwater supplied from the Buena Vista Water Storage District (BVWSD) located to the northwest. The raw water supply pipeline will be approximately 18 miles in length. Potable water for drinking and sanitary use will be supplied by West Kern Water District located near the State Route 119 (SR 119)/Tupman Road intersection (southeast of the Project Site). The potable water supply pipeline will be approximately 5.5 miles in length.
- **Carbon Dioxide Pipeline** – The carbon dioxide pipeline will transfer the carbon dioxide captured during gasification from the Project Site southwest to the custody transfer point. The Project is considering two alternative pipeline routes. Alternative 1 is approximately 2 miles in length, while Alternative 2 is approximately 2.5 miles in length.

The Project components described above are shown on Figure 2-5, Project Location Map, which depicts the region, the vicinity, the Project Site and its immediate surroundings for Project components.

This section of the AFC summarizes the potential environmental impacts on paleontological resources that could result from construction of the Project. This paleontological resources inventory and impact assessment was prepared by Dr. Lanny H. Fisk, PhD PG, a California licensed Professional Geologist (PG) and Principal Paleontologist, and by Stephen J. Blakely, Project Manager and Staff Paleontologist, both with PaleoResource Consultants (PRC). It meets all requirements of the California Energy Commission (CEC) (CEC 2007) and the standard measures for mitigating adverse construction-related environmental impacts on significant

paleontological resources established by the Society of Vertebrate Paleontology (SVP 1995, 1996).

Paleontological resources (fossils) are the remains or traces of prehistoric animals and plants. Fossils are important scientific and educational resources because of their use in (1) documenting the presence and evolutionary history of particular groups of now extinct organisms, (2) reconstructing the environments in which these organisms lived, (3) determining the relative ages of the strata in which they occur, and (4) determining the geologic events that resulted in the deposition of the sediments in which they were buried.

### 5.16.1 Affected Environment

#### *5.16.1.1 Geographic Location*

The Project Site is located within 1 mile (approximately [~] 1.6 kilometers) of the border of the Elk Hills Oil and Gas Field (Figure 2-5, Project Location Map) within Section 22 of Township 30 South, Range 24 East. The center of the Project Site is at approximately latitude 35°18'28" North and longitude 119°23'08" West.

The Project will impact Section 22 within Township 30 South, Range 24 East on the East Elk Hills U.S. Geological Survey (USGS) 7.5-minute Quadrangle. The water pipeline will impact the following:

- Sections 27 and 34 within Township 28 South, Range 22 East;
- Sections 1, 2, and 12 within Township 29 South, Range 22 East;
- Sections 7, 17, 18, 20, 21, 27, 28, and 34 through 36 within Township 29 South, Range 23 East;
- Section 1 within Township 30 South, Range 23 East; and
- Sections 5, 6, 8 through 10, 14 through 16, 22, and 23 within Township 30 South, Range 24 East.

The carbon dioxide pipeline (considering two alternatives) will impact Sections 21, 22, 28, and 33 within Township 30 South, Range 24 East. The natural gas pipeline will impact the following:

- Sections 22 through 25 within Township 30 South, Range 24 East;
- Sections 30 through 33 within Township 30 South, Range 25 East; and
- Sections 4 and 5 within Township 31 South, Range 25 East.

The transmission line (considering two alternatives) will impact the following:

- Section 13 within Township 29 South, Range 23 East;
- Sections 18 through 21 and 28 through 33 within Township 29 South, Range 24 East; and
- Sections 4 through 9, 16 and 17, and 21 and 22 within Township 30 South, Range 24 East.

The Project area is located at the north edge of the Elk Hills, which are near the western border of the San Joaquin Valley. The San Joaquin Valley comprises roughly the southern two-thirds of

the major north-northwest-oriented synclinorium called either the Valle Grande (Clark 1929), Great Valley (Fenneman 1931; Hackel 1966), Great Interior Valley (Harradine 1950), Great San Joaquin Valley (Piper et al. 1939; Davis et al. 1957), or California Trough (Piper et al. 1939). The Great Valley Physiographic Province is located between the Sierra Nevada Physiographic Province on the east and the Coast Ranges Physiographic Province on the west (Jahns 1954).

The Project Site is located near the northern margin of the Elk Hills, which are approximately 17 miles (~27 kilometers) long and 7 miles (~11 kilometers) wide. The Elk Hills reach an elevation of 1,551 feet (~473 meters), which is approximately 1,200 feet (~366 meters) above the floor of the San Joaquin Valley (Berryman 1973). Access to the Project Site is provided from Tupman Road.

### 5.16.1.2 Regional Geologic Setting

The general geology of the San Joaquin Valley has been described in some detail by Mendenhall (1908), Mendenhall et al. (1916), Piper et al. (1939), Hoots et al. (1954), Davis et al. (1957, 1959, 1964), Davis and Hall (1959), Hoffman (1964), Croft and Wahrhaftig (1965), Hackel (1966), Croft and Gordon (1968), Bull (1973), Page (1986), Marchand (1977), Bartow and Marchand (1979), Marchand and Allwardt (1981), Lettis (1988), Bartow (1987, 1991), Beyer and Bartow (1988), Callaway and Rennie (1991), and Lettis and Unruh (1991), among others.

Only a few authors have specifically described the geology in the vicinity of the unincorporated community of Tupman or the Elk Hills Oil Field Unit, including Woodring et al. (1932), Porter (1943), Wells (1952), Adkison (1973), Berryman (1973), Dibblee (1973), and Maher et al. (1975). Surficial geologic mapping of the Project vicinity has been provided at a scale of 1:1,000,000 by Wahrhaftig et al. (1993); at a scale of 1:750,000 by Jennings (1977); at a scale of 1:500,000 by Mendenhall et al. (1916), Jenkins (1938), and Bartow (1987, 1991); at a scale of approximately 1:320,000 by Morton and Troxel (1962); at a scale of 1:250,000 by Smith (1964); at a scale of 1:62,500 by Dibblee (1972); at a scale of 1:31,680 by Woodring et al. (1932); and at a scale of 1:24,000 by Dibblee (2005a, 2005b, 2005c, 2005d, 2005e, 2005f).

The information in these geologic maps and published and unpublished reports form the basis of the following discussion. Individual maps and publications are incorporated into this report and referenced where appropriate. For obtaining the older geological literature, the exhaustive compilation entitled "Geological Literature on the San Joaquin Valley of California" by Maher et al. (1973) was particularly helpful. The aspects of geology pertinent to this report are the types, distribution, and age of sediments immediately underlying the Project area and their probability of producing fossils during construction. The site-specific geology in the vicinity of the Project is discussed separately below.

The San Joaquin Valley is a great structural depression between the westerly tilted Sierra Nevada block on the east and the complexly folded and faulted Coast Ranges on the west. The valley is filled with thick Mesozoic and Tertiary marine and non-marine sediments covered by a relatively thin veneer of Quaternary alluvial sediments (Bailey 1966). The Elk Hills are located along the western edge of the San Joaquin Valley, where they rise above the surrounding relatively flat valley. The Elk Hills are the topographic expression of the Elk Hills Anticline, which is part of the *en echelon* folding of the Tertiary and Quaternary sedimentary strata along the western side of the San Joaquin Valley (White 1987). The axes of these folds trend generally northwest-

southeast, and are associated with strain caused by movement along the San Andreas Fault (White 1987).

### *5.16.1.3 Resource Inventory Methods*

To develop a baseline paleontological resource inventory of the Project Site and surrounding area and to assess the potential paleontological productivity of each stratigraphic unit present, the published as well as available unpublished geological and paleontological literature was reviewed. Stratigraphic and paleontologic inventories were compiled, synthesized, and evaluated (see below). These methods are consistent with CEC (2007) and Society of Vertebrate Paleontology (SVP) (1995) guidelines for assessing the importance of paleontological resources in areas of potential environmental effect.

Geologic maps and reports covering the bedrock and surficial geology of the Project vicinity were reviewed to determine the exposed and subsurface rock units, to assess the potential paleontological productivity of each rock unit, and to delineate their respective areal distribution in the Project area. Museum records searches were conducted at the University of California Museum of Paleontology (UCMP) at Berkeley, the Los Angeles County Natural History Museum (LACM), and the San Bernardino County Museum of Natural History (SBMNH) in order to determine whether any of the stratigraphic units found within the Project vicinity had previously yielded significant paleontological resources. In addition, aerial photographs of the area were examined to aid in determining the areal distribution of distinctive sediment and soil types. No subsurface exploration was conducted for this assessment. However, a PRC field paleontologist was present during augering for geotechnical boreholes on the Project Site, and did observe subsurface stratigraphy and fossils (see discussion below).

A field survey, which included visual inspection of exposures of potentially fossiliferous strata in the Project area, was conducted to document the presence of sediments suitable for containing fossil remains and the presence of any previously unrecorded fossil sites. The field survey for this assessment was conducted over several site visits from March 2008 through May 2008. Dr. Lanny H. Fisk, PhD, PG is the principal paleontologist with PRC and surveyed during 04 and 12 March and 14-15 and 20-21 May. Dr. Hugh M. Wagner, PhD is a senior paleontologist with PRC and surveyed during 04-08 and 09-12 March and during 29 April. Mr. Patrick W. Riseley, PG is a field paleontologist with PRC and surveyed 02-07, 09-13, 19-21, and 31 March, 01-04 and 06-07 April, and 20-22 May. David M. Maloney, field supervisor with PRC, surveyed the site 05-06 March and 20-22 May. John N. Adrian, field paleontologist with PRC, surveyed 31 March and 01-04 April. Phil R. Peck, field paleontologist with PRC, surveyed 08-09, 12-15, 20-23, and 28 May. Richard J. Serrano, field paleontologist with PRC, surveyed 08-10, 12-15, 20-24, and 28-29 May. During the field survey, stratigraphy was observed in arroyos, hill-slopes, badlands, and road cuts. Exposed sediments up to approximately 30 feet (~9 meters) were observed in locations in the vicinity of the Project.

### *5.16.1.4 Paleontological Resource Assessment Criteria*

The SVP (1995), in common with other environmental disciplines such as archaeology and biology (specifically in regard to listed species), considers any fossil specimen significant unless demonstrated otherwise, and protected by environmental statutes. This position is held because fossils are uncommon and only rarely will a fossil locality yield a statistically significant number

of specimens representing the same species. In fact, vertebrate fossils are so uncommon that, in most cases, each fossil specimen found will provide additional important information about the characteristics or distribution of the species it represents.

A stratigraphic unit (such as a formation, member, or bed) known to contain significant fossils is considered to be “sensitive” to adverse impacts if there is a high probability that earth-moving or ground-disturbing activities in that rock unit will either disturb or destroy fossil remains. This definition of sensitivity differs fundamentally from that for archaeological resources:

*It is extremely important to distinguish between archaeological and paleontological (fossil) resource sites when defining the sensitivity of rock units. The boundaries of archaeological sites define the areal extent of the resource. Paleontologic sites, however, indicate that the containing sedimentary rock unit or formation is fossiliferous. The limits of the entire rock formation, both areal and stratigraphic, therefore define the scope of the paleontologic potential in each case (SVP 1995).*

This distinction between archaeological and paleontological sites is important. Most archaeological sites have a surface expression that allow for their geographic location. Fossils, on the other hand, are an integral component of the rock unit below the ground surface; therefore, they are not observable unless exposed by erosion or human activity. Thus, a paleontologist cannot know either the quality or quantity of fossils present before the rock unit is exposed as a result of natural erosion processes or earth-moving activities. The paleontologist can only make conclusions on sensitivity to impact based upon what fossils have been found in the rock unit in the past, along with a judgment on whether or not the depositional environment of the sediments that compose the rock unit was likely to result in the burial and preservation of fossils.

Fossils are seldom uniformly distributed within a rock unit. Most of a rock unit may lack fossils, but at other locations within the same rock unit concentrations of fossils may exist. Even within a fossiliferous portion of the rock unit, fossils may occur in local concentrations. For example, Shipman (1977, 1981) excavated a fossiliferous site using a three-dimensional grid and removed blocks of matrix of a consistent size. The site chosen was known prior to excavation to be richly fossiliferous, yet only 17 percent of the blocks actually contained fossils. These studies demonstrate the physical basis for the difficulty in predicting the location and quantity of fossils in advance of project-related ground disturbance.

Since it is unfortunately not possible to determine where fossils are located without actually disturbing a rock unit, monitoring of excavations by an experienced paleontologist during construction increases the probability that fossils will be discovered and preserved. Preconstruction mitigation measures, such as surface prospecting and collecting, will not prevent adverse impacts on fossils because many sites will be unknown in advance due to an absence of fossils at the surface.

The non-uniform distribution of fossils within a rock unit is typical. Many paleontological resource assessment and mitigation reports conducted in support of environmental impact documents and mitigation plan summary reports document similar findings (see for instance Lander 1989, 1993; Reynolds 1987, 1990; Spencer 1990; Fisk et al. 1994; and references cited therein). In fact, most fossil sites recorded in reports of impact mitigation (where construction monitoring has been implemented) had no previous surface expression. Because the presence or

location of fossils within a rock unit cannot be known without exposure resulting from erosion or excavation, under SVP (1995) standard guidelines, an entire rock unit is assigned the same level of sensitivity based on recorded fossil occurrences.

Using SVP (1995) criteria, the paleontological importance or sensitivity (high, low, or undetermined) of each rock unit exposed in a project site or surrounding area is the measure most amenable to assessing the significance of paleontological resources because the areal distribution of each rock unit can be delineated on a topographic or geologic map. The paleontological sensitivity of a stratigraphic unit reflects: (1) its potential paleontological productivity, and (2) the scientific significance of the fossils it has produced. This method of paleontological resources assessment is the most appropriate because discrete levels of paleontological importance can be delineated on a topographic or geologic map.

The potential paleontological productivity of a stratigraphic unit exposed in a project area is based on the abundance/densities of fossil specimens and/or previously recorded fossil sites in exposures of the unit in and near a project site. The underlying assumption of this assessment method is that exposures of a stratigraphic unit in a project site are most likely to yield fossil remains both in quantity and density similar to those previously recorded from that stratigraphic unit in and near the project site.

Under California Environmental Quality Act of 1970 (CEQA) Guidelines §15064.5 (a) (2), public agencies must treat all historical and cultural resources as significant unless the preponderance of evidence demonstrates that they are not historically or culturally significant. An individual fossil specimen is considered scientifically important if it is:

- Identifiable
- Complete
- Well preserved
- Age diagnostic
- Useful in paleo-environmental reconstruction
- A type or topotypic specimen
- A member of a rare species
- A species that is part of a diverse assemblage
- A skeletal element different from, or a specimen more complete than, those now available for that species

All identifiable land mammal fossils are considered scientifically important because of their potential use in providing relative age determinations and paleo-environmental reconstructions for the sediments in which they occur. Moreover, vertebrate remains are comparatively rare in the fossil record. Although fossil plants are usually considered of lesser importance because they are less helpful in age determination, they are actually more sensitive indicators of their environment (Miller et al. 1971) and as sedentary organisms, are more valuable than mobile animals for paleo-environmental reconstructions. For marine sediments, invertebrate and marine algal fossils, including microfossils, are scientifically important for the same reasons that land mammal and/or land plant fossils are valuable in terrestrial deposits. The value or importance of

different fossil groups varies depending on the age and depositional environment of the stratigraphic unit that contains the fossils.

The following tasks were completed to establish the paleontological importance and sensitivity of each stratigraphic unit exposed in or near the Project Site:

- The potential paleontological productivity of each rock unit was assessed based on previously recorded and newly documented fossil sites it contains at and/or near the Project Site.
- The scientific importance of fossil remains recorded from a stratigraphic unit exposed at and/or near the Project Site were assessed.
- The paleontological importance of a rock unit was assessed, based on its documented and/or potential fossil content in the area surrounding the Project Site.

### **Categories of Sensitivity**

In its standard guidelines for assessment and mitigation of adverse impacts to paleontological resources, the SVP (1995) established three categories of sensitivity for paleontological resources: high, low, and undetermined.

**High Sensitivity.** Stratigraphic units in which fossils have been previously found have a high potential to produce additional fossils and are therefore considered to be highly sensitive. In the significance criteria of the SVP (1995), all vertebrate fossils are categorized as having significant scientific value and all stratigraphic units in which vertebrate fossils have previously been found have high sensitivity. In areas of high sensitivity, full-time monitoring is recommended during any project-related ground disturbance.

**Low Sensitivity.** Stratigraphic units that are not sedimentary in origin or that have not been known to produce fossils in the past are considered to have low sensitivity. Monitoring is usually not recommended nor needed during excavation in a stratigraphic unit with low sensitivity.

**Undetermined Sensitivity.** Stratigraphic units that have not had any previous paleontological resource surveys or any fossil finds are considered to have undetermined sensitivity. After reconnaissance surveys, observation of artificial exposures (e.g., road cuts) and natural exposures (e.g., stream banks), and possible subsurface testing (e.g., augering or trenching), an experienced, professional paleontologist can often determine whether the stratigraphic unit should be categorized as having high or low sensitivity.

The Bureau of Land Management (BLM) Paleontological Resources Handbook H-8270-1 (BLM 1998) uses a slightly different classification system for ranking areas according to their potential to contain significant fossils. These rankings are used in land use planning, as well as to identify areas that may warrant special management and/or special designation such as Areas of Critical Environmental Concern. Public lands managed by BLM are classified based on their potential to contain fossils using the following criteria:

Condition 1 – Areas that are known to contain vertebrate fossils or noteworthy occurrences of invertebrate or plant fossils.

Condition 2 – Areas with exposures of geological units or settings that have high potential to contain vertebrate fossils or noteworthy occurrences of invertebrate or plant fossils.

Condition 3 – Areas that are very unlikely to produce vertebrate fossils or noteworthy occurrences of invertebrate or plant fossils based on their surficial geology, igneous or metamorphic rocks, extremely young alluvium, colluvium, or aeolian deposits, or the presence of deep soils (BLM 1998).

A 1971 BLM (Marshall 1976, p. 347) internal memorandum suggested the following criteria for determining the significance of individual paleontological resources. “A paleontological resource may be significant if:

- It represents a rare species or one that has not been recorded previously in the literature.
- It illustrates previously unknown sexual dimorphism, phenotypic variation, or an ontogenetic series of a given taxon.
- It is from a locality that marks either a geographical or temporal range extension for a given species.
- It is exceptional in that it represents an exhibit-quality specimen.
- It represents material that assists in refining the age assignment of an otherwise poorly dated litho-stratigraphic unit.
- It represents a concentration of vertebrate specimens in a bed or series of beds. The sample may include either associated skeletal material referable to an individual or an aggregate of specimens referable to more than one individual. In either case, the material yields potentially significant taphonomic information that can be utilized in paleontologic analyses.
- It provides important information on the evolutionary trends among organisms, relating inhabitants of the earth to extinct organisms.
- It provides important information regarding development of biological communities or interaction between botanical and zoological biota.
- It demonstrates unusual or spectacular circumstances in the history of life.
- It is in short supply and in danger of being depleted or destroyed by the elements, vandalism, or commercial exploitation, and is not found in other geographic localities.
- All vertebrate fossils are of scientific value.”

The previously described BLM criteria have been widely used by both lead agencies and professional mitigation paleontologists as objective measures of significance. In this paleontological resource impact assessment, the criteria of both the SVP (1995) and the BLM (1998) are applied. BLM lands will not be directly impacted by the Project or by any of its associated linears. However, BLM lands do occur within the 1-mile paleontological survey area, so the criteria will be considered.

### 5.16.1.5 Resource Inventory Results

#### Stratigraphic Inventory

Regional geologic mapping in the vicinity of the Project Site has been provided by Jennings (1977; 1:750,000), Mendenhall et al. (1916; 1:500,000), Jenkins (1938; 1:500,000), Bartow (1987, 1991; 1:500,000), Morton and Troxel (1962; ~1:320,000), and Smith (1964; 1:250,000). Larger scale mapping of the Project Site has been provided by Dibblee (1972; 1:62,500), Woodring et al. (1932; 1:31,680), and Dibblee (2005a-2005f; 1:24,000).

#### Project Geology

Based upon the available geologic literature, recent geologic maps, and field observations, two stratigraphic units will be potentially impacted during Project construction activities. In the discussion below, the stratigraphic nomenclature of the area will follow that of Dibblee (2005a through 2005f), the most detailed and also most recent geologic maps available. Dibblee (2005a through 2005f) identified two stratigraphic units within the Project vicinity: Quaternary alluvium and Tulare Formation. Each of these stratigraphic units is described below.

In his geologic mapping, Dibblee (2005a through f) mapped the area in the vicinity of the Project Site and the ROWs of the linears as either Quaternary alluvium or Tulare Formation. The Project Site is mapped as Quaternary alluvium, although the map indicates that this alluvium unconformably overlies sediments of the Tulare Formation (Dibblee 2005f). Thus, although Quaternary alluvium is mapped as being present at the surface over the Project Site, the older Tulare Formation may still be encountered in the shallow subsurface. This was confirmed through the observation of augering for geotechnical boreholes when an unconformity consistent with Quaternary alluvium overlying sediments of Tulare Formation was observed at approximately 4.5 feet (~ 1.4 meters) below ground surface. Linear facilities associated with Project construction will also potentially impact sediments of the Tulare Formation. Many of the linear facilities will at some point pass through areas mapped as Tulare Formation or areas mapped as Quaternary alluvium overlying Tulare Formation (Dibblee 2005a through 2005f).

Excavations at the Project Site and along the ROWs of the water pipeline and electrical transmission lines will disturb sediments of Quaternary alluvium, and potentially Tulare Formation. Excavations along the ROWs of the natural gas and CO<sub>2</sub> pipelines will disturb sediments of both the Quaternary alluvium and the Plio-Pleistocene Tulare Formation.

**Tulare Formation.** Late Pliocene to Pleistocene age Tulare Formation was named by Anderson (1905), who did not designate a type section. Woodring et al. (1940) later designated the Kettleman Hills North Dome as the type section for the formation. Dibblee (1973) described the Tulare Formation as “locally deformed dissected valley deposits composed of gravel, sand, and silt.” Lithologically, the Tulare Formation consists of argillaceous sand and silt deposits with lenses of coarse sand and gravel. White (1987) described sediments of the Tulare Formation as found in the Elk Hills as “low-angle, cross-bedded, fine to medium pebbly sands interbedded with structureless to faintly laminated, gypsiferous, olive-green, brown and gray muds and clays. Conglomerate units do occur, but are rare overall. Pebbles and clasts of siliceous shale are common and are most likely derived from the Monterey Formation exposed in the Temblor Range to the west.” Tulare Formation sediments in the Elk Hills have a thickness of up to approximately 610 meters (~2,000 feet), while Tulare sediments found elsewhere may be as

much as 1,525 meters (~5,000 feet) thick (Maher et al. 1975; White 1987). Most of the formation is composed of reworked sedimentary materials whose origin is from erosion of the Coast Ranges. The Tulare Formation overlies the San Joaquin Formation, likely conformably, in the Elk Hills area, though in other places throughout the San Joaquin Valley it unconformably overlies sediments of various formations and ages (Dibblee 1973; Lettis 1982). The age of the Tulare Formation has been determined based upon structural and stratigraphic relationships, paleontological correlations, radiometric dating methods, and paleomagnetic data. White (1987) used measured magnetic polarities within the Tulare Formation from locations in the southern San Joaquin Valley to determine this age to be between 0.90 million years and 2.48 million years.

**Quaternary Alluvium.** Quaternary alluvium is composed primarily of fluvial sands and gravels reworked from older formations and transported from the topographically-high adjacent areas. Within and in the immediate vicinity of the Project Site, the alluvium is primarily composed of either reworked Tulare Formation material and recent soils, or sediments of the Kern River distal fan. There is also some lacustrine material in the local alluvium, including sediments of Buena Vista Lake and other periodic lakes. Two drill sites located northeast of Buttonwillow, produced fossil wood which was analyzed using radiometric dating methods (Manning 1968). These samples, recovered at 20 and 35 feet below ground surface, produced a late Pleistocene age.

### **Paleontological Resource Inventory**

An inventory of known paleontological resources previously discovered in the vicinity of the Project is presented below and the paleontological importance of these resources is assessed. The literature review and UCMP, LACM, and SBMNH archival records search conducted for this inventory documented no previously recorded fossil sites within the actual Project Site. Previously reported fossil sites do occur within 1 mile of Project Site and linear facilities, and numerous previously unreported fossil sites were identified during the field survey for this Project. In addition, sediments of Quaternary alluvium and Plio-Pleistocene Tulare Formation have yielded fossilized remains of extinct species of continental vertebrates and other types of organisms at previously recorded fossil sites in the region (Jefferson 1991a, 1991b; UCMP records; others described below).

**Tulare Formation.** The Tulare Formation has yielded fossil remains at numerous sites in the San Joaquin Valley. These remains include algal stromatolites (vertically layered mat-like algal growths); diatoms; petrified wood; shells of snails and clams; and the bones and teeth of bony fishes, amphibians, turtles, lizards, snakes, birds, and a diversity of extinct land mammals, including moles, ground sloths, rabbits, squirrels, gophers, pocket mice, kangaroo rats, pack rats, deer mice, cotton rats, grasshopper mice, dogs, saber-tooth cats, horses, peccaries, camels, tapirs, and deer (Anderson and Pack 1915; Arnold and Johnson 1910; Davis et al. 1957, 1959; Foss and Blaisdell 1968; Gester 1917; Hoots et al. 1954; Lander 1993; Maher et al. 1975; Merriam 1903, 1905, 1914, 1915a, 1915b, 1917; Porter 1943; Repenning 1980; Reynolds 1987, 1990; Stirton and VanderHoof 1933; Taylor 1966; Wood and Davis 1959; Woodring et al. 1932; UCMP records). Anderson and Pack (1915) also mentioned recycled fossils from older stratigraphic units and silicified wood in the Tulare Formation.

There are a number of previously recorded fossil sites in the Tulare Formation within approximately 15 miles of the Project Site. These include several sites in the Elk Hills in addition to several others in neighboring areas such as McKittrick (Woodring et al. 1932;

Jefferson 1991a, 1991b; UCMP records). Several fossil localities described by Woodring et al. (1932) are present in the Elk Hills, and include specimens of camel, horse, rabbit, wood rat, cotton rat, and silicified wood. Additionally, Woodring et al. (1932) described freshwater invertebrates from “ditch samples” in the Elk Hills. Based upon these fossil localities, Woodring et al. (1932) stated that “the Elk Hills offer a promising field for collecting vertebrate fossils, which would fill a gap in the succession of vertebrate faunas on the Pacific coast.” Maher et al. (1975) indicated that “scattered fish remains,” mollusk fragments, reworked foraminifers, ostracodes, pelecypods, and small gastropods had been identified from wells in the Elk Hills. Jefferson (1991a, 1991b) in his previously described database of California Late Pleistocene (Rancholabrean North American Land Mammal Age [NALMA]) vertebrate fossils, indicated two localities in the Tulare Formation.

During the field survey for prospective fossil localities, many previously unrecorded sites were identified as occurring within 1 mile of the Project Site and its associated linear facilities. Fossils at these localities included vertebrate fossil bones and bone fragments, invertebrate shells, and fossilized wood. Numerous paleosols were also identified within the Tulare Formation which contained ichnofossils.

In summary, sediments referable to the Tulare Formation have yielded an abundance of invertebrate, vertebrate, and plant fossils, including microfossils. Several previously recorded and previously unrecorded fossil localities are found in the area of the Project Site, including several sites within the Elk Hills (Jefferson 1991b; UCMP records). Because this unit has in the past produced significant vertebrate fossils, the Tulare Formation is categorized as highly sensitive (SVP 1995).

***Quaternary Alluvium.*** No fossil localities have previously been reported from Quaternary alluvium at the Project Site. However, significant vertebrate fossils have been reported from Holocene and Pleistocene sediments in several areas of Kern County (Jefferson 1991a, 1991b; UCMP records). Jefferson (1991a, 1991b) compiled a database of California Late Pleistocene (Rancholabrean NALMA) to earliest Holocene vertebrate fossils from published records, technical reports, unpublished manuscripts, information from colleagues, and inspection of museum paleontological collections at over 40 public and private institutions. He listed over 70 individual sites in Kern County that yielded vertebrate fossils of these ages. Many of these sites are not assigned to a specific formation, group, or member, and may be referable to sediment of unnamed (Quaternary) alluvium. Among these localities is a Rancholabrean vertebrate fossil locality discovered during construction of the Bakersfield Canal (UCMP V-65247). Fossils discovered during this construction project have been identified as an extinct species of horse. Additionally, Pleistocene fossil wood was recovered from well borings 10 to 15 miles northwest of Buttonwillow (Manning 1968). During the field survey performed for this Project, previously unrecorded fossil localities were identified. Specimens identified at these sites included freshwater invertebrate shells and ichnofossils. Fossils occurring in Quaternary alluvium are valuable to the scientific community as they provide information about climatic conditions in the not too distant past. The occurrence of large and small mammals are well documented from these and older subsurface deposits and with further observation of earth-moving activities and prospecting for fossils, more specimens could be unearthed. Since fossil vertebrates have been previously reported from Quaternary alluvium within Kern County, the Quaternary alluvium is also judged to have high sensitivity based on SVP guidelines (1995).

**Summary**

In summary, although no fossils were previously reported to directly underlie the Project, numerous fossil localities nearby within the Quaternary alluvium and the Tulare Formation have been reported in both the published scientific literature and museum records. In addition, numerous previously unrecorded fossil localities were identified during the field survey of the Project and linear facility ROWs. Many of these previously reported and unreported localities occur within 1 mile of Project related features. The presence of fossils in sediments of Quaternary alluvium within one mile of the Project and elsewhere in Kern County, and of fossils in sediments of Plio-Pleistocene Tulare Formation within one mile of the Project, elsewhere, and in the Elk Hills suggests that there is a high potential for additional similar fossil remains to be uncovered by excavations during Project construction. Under SVP (1995) criteria, these stratigraphic units have a high potential for producing additional sensitive paleontological resources.

**5.16.2 Environmental Consequences**

Potential impacts on paleontological resources resulting from construction of the Project can be divided into construction-related impacts and operation-related impacts. The potential environmental effects from construction and operation of the Project on paleontological resources are presented in the following subsections.

**Potential Impacts from Project Construction.** Construction-related impacts to paleontological resources primarily involve terrain modifications (excavations and drainage diversion measures). Paleontological resources, including an undetermined number of fossil remains and unrecorded fossil sites; associated specimen data and corresponding geologic and geographic site data; and the fossil-bearing strata, could be adversely affected by (i.e., will be sensitive to) ground disturbance and earth moving associated with construction of the Project. Direct impacts could result from: vegetation clearing; grading of roads, drainage diversions, pile driving, and other ground disturbance at the Project Site; trenching for pipelines; augering for foundations for electrical transmission towers or poles; and any other Project-related earth-moving activity.

Identifiable fossil remains recovered from the Quaternary alluvium or the Plio-Pleistocene Tulare Formation during Project construction could be scientifically important and significant. These fossil remains discovered during construction could represent new taxa or new fossil records for the San Joaquin Valley, for the state of California, for the Quaternary/Tertiary eras, or for a stratigraphic unit. They could also represent geographic or temporal range extensions. Moreover, discovered fossil remains could make it possible to more accurately determine the age, paleo-climate, and depositional environment of the sediments from which they are salvaged. Finally, fossil remains salvaged during Project construction could provide a more comprehensive documentation of the diversity of animal and plant life that once existed in Kern County and could result in a more accurate reconstruction of the geologic and paleo-biologic history of the San Joaquin Valley and the Elk Hills area.

Clearing, grading, and excavations that encounter previously undisturbed sediment at the Project could result in significant adverse impacts to paleontological resources. These construction impacts could disturb previously undisturbed fossiliferous sediments, that without mitigation, could make those sediments and their paleontologic resources unavailable for future scientific

investigation. However, with the properly designed and implemented mitigation program, these impacts will be reduced to less than significant.

**Potential Impacts from Project Operation.** No impacts on paleontological resources are expected to occur from the continuing operation of the Project or any of its related facilities.

### 5.16.3 Cumulative Impacts Analyses

If paleontological finds were to be encountered during Project construction, the potential for cumulative impacts will exist. Mitigation measures will be implemented to salvage such resources and reduce cumulative impacts to a level that is less than significant. The mitigation measures described in Section 5.16.4, Mitigation Measures, will effectively preserve the value to science of any significant fossils uncovered during Project-related construction.

### 5.16.4 Mitigation Measures

This section describes mitigation measures that will be implemented to reduce potential adverse impacts to significant paleontological resources resulting from Project construction. Mitigation measures are necessary because of potential adverse impacts of Project construction on significant paleontological resources within the Quaternary alluvium, and the Plio-Pleistocene Tulare Formation. The paleontological resource impact mitigation program will reduce to an insignificant level the direct, indirect, and cumulative adverse environmental impacts on paleontological resources that could result from Project construction. The mitigation measures summarized below are consistent with SVP standard guidelines for mitigating adverse construction-related impacts on paleontological resources (SVP, 1995, 1996), and fulfill the requirements of the BLM (1998).

Implementation of these mitigation measures will reduce the potentially significant adverse environmental impact of Project-related ground disturbance and earth-moving on paleontological resources to an insignificant level by allowing for the salvage of fossil remains and associated specimen data and corresponding geologic and geographic site data that otherwise might be lost to earth-moving and to unauthorized fossil collecting.

With a well designed and implemented paleontological resource monitoring and mitigation plan, Project construction could actually result in beneficial impacts on paleontological resources through the discovery of fossil remains that would not have been exposed without Project construction and, therefore, would not have been available for study. The salvage of fossil remains as part of Project construction could help answer important questions regarding the geographic distribution, stratigraphic position, and age of fossiliferous sediments in the Project area.

#### **PALEO-1 Paleontological Monitoring.**

Prior to construction, a qualified paleontologist will be retained to both design and implement a monitoring and mitigation program. During construction, ground-disturbing activities will be monitored where these activities will potentially disturb previously undisturbed sediment. Monitoring will not be conducted in areas where the ground has been previously disturbed or in areas where exposed sediment will be buried, but not otherwise disturbed. Construction monitoring will be conducted to ensure that unanticipated discoveries are addressed in a timely manner.

**PALEO-2 Paleontological Monitoring and Mitigation Program.**

The paleontological resource monitoring and mitigation program will include preconstruction coordination; construction monitoring; emergency discovery procedures; sampling and data recovery, if needed; preparation, identification, analysis, and museum curation of any fossil specimens and data recovered; and reporting. This monitoring and mitigation plan will be consistent with SVP (1995) standard guidelines for the mitigation of construction-related adverse impacts on paleontological resources, as well as the requirements of the designated museum repository for any fossils collected (SVP 1996).

**PALEO-3 Construction Personnel Education.**

Prior to start of Project construction, construction personnel involved with earth-moving activities should be informed: (1) that fossils may be discovered during excavating, (2) that these fossils are protected by laws, (3) on the appearance of common fossils, and, (4) on proper notification procedures. This worker training should be prepared and presented by a qualified paleontologist.

**PALEO-4 Paleontological Monitoring.**

Prior to the start of construction, the paleontologist should conduct a field survey of exposures of sensitive stratigraphic units that will be disturbed and any fossils discovered should be salvaged. Earth-moving construction activities should be monitored wherever these activities will disturb previously undisturbed sediment. Monitoring will not need to be conducted in areas where sediments have been previously disturbed or in areas where exposed sediments will be buried, but not otherwise disturbed.

**5.16.5 Laws, Ordinances, Regulations, and Standards**

Paleontological resources are classified as non-renewable scientific resources and are protected by several federal and state statutes (California Office of Historic Preservation 1983; Marshall 1976; West 1991; Fisk and Spencer 1994; Gastaldo 1999), most notably by the 1906 Federal Antiquities Act and other subsequent federal legislation and policies and by the state of California's environmental regulations (CEQA, §15064.5). Professional standards for assessment and mitigation of adverse impacts on paleontological resources have been established by the SVP (1995, 1996). Design, construction, and operation of the Project, including ancillary facilities, will be conducted in accordance with laws, ordinances, regulations, and standards (LORS) applicable to paleontological resources. Federal and state LORS applicable to paleontological resources are summarized in Table 5.16-1, Summary of LORS – Paleontological Resources, and discussed briefly below, together with county and city requirements and SVP professional standards.

**Table 5.16-1  
Summary of LORS – Paleontological Resources**

| <b>LORS</b>                           | <b>Requirements</b>                                 | <b>Conformance Section</b> | <b>Administering Agency</b>     | <b>Agency Contact</b>           |
|---------------------------------------|---|----------------------------|---------------------------------|---------------------------------|
| <b>State Jurisdiction</b>             |   |                            |                                 |                                 |
| CEQA                                  | Protects paleontological resources on state lands.  | 5.16.5                     | CEC                             | Eileen Allen<br>916-654-4082    |
| Public Resources Code §5097.5/§5097.9 | Protects paleontological resources on state lands.  | 5.16.5                     | CEC                             | Eileen Allen<br>916-654-4082    |
| <b>Local Jurisdiction</b>             |   |                            |                                 |                                 |
| Kern County General Plan              | Protects paleontological resources on county lands. | 5.16.5                     | Kern County Planning Department | Cheryl Casdorff<br>661-862-8600 |

Source: HECA Project

Notes:

CEC = California Energy Commission

CEQA = California Environmental Quality Act

LORS = laws, ordinances, regulations, and standards

**5.16.5.1 Federal**

There are no applicable federal LORS because the Project affects no federal lands.

**5.16.5.2 State**

The CEQA lead agency having jurisdiction over a project is responsible to ensure that paleontological resources are protected in compliance with CEQA and other applicable statutes. California Public Resources Code § 21081.6, entitled Mitigation Monitoring Compliance and Reporting, requires that the lead agency demonstrate project compliance with mitigation measures developed during the environmental impact review process.

Other state requirements for paleontological resources management are in Public Resources Code Chapter 1.7, § 5097.5 (Stats. 1965, c. 1136, p. 2792), entitled Archaeological, Paleontological, and Historical Sites. This statute defines any unauthorized disturbance or removal of a fossil site or fossil remains on public land as a misdemeanor and specifies that state agencies may undertake surveys, excavations, or other operations as necessary on state lands to preserve or record paleontological resources. This statute will apply to the Project if the Project will be built on city-owned or state managed lands.

**5.16.5.3 Local**

California Planning and Zoning Law requires each county and city jurisdiction to adopt a comprehensive, long-term general plan for its development. The general plan is a policy document designed to give long-range guidance to those making decisions affecting the future character of the planning area. It represents the official statement of the community’s physical

development as well as its environmental goals. The general plan also acts to clarify and articulate the relationship and intentions of local government to the rights and expectations of the general public, property owners, and prospective investors. Through its general plan, the local jurisdiction informs these groups of its goals, policies, and development standards; thereby, communicating what must be done to meet the objectives of the general plan. State planning law requires each jurisdiction to identify environmental resources and to prepare and implement policies which relate to the utilization and management of these resources.

The Kern County General Plan addresses paleontological resources in the Land Use, Open Space, and Conservation Element under “General Provisions 1.10.3: Archaeological, Paleontological, Cultural, and Historical Preservation.” Under this heading, Policy 25 states, “the County will promote the preservation of cultural and historic resources which provide ties with the past and constitute a heritage value to residents and visitors.” Implementation Measure L. for this Policy states that “the County shall address archaeological and historical resources for discretionary projects in accordance with CEQA.” Implementation Measure M. for this Policy states that “in areas of known paleontological resources, the County should address the preservation of these resources where feasible.”

#### **5.16.6 Involved Agencies and Agency Contacts**

Other than CEC, no state or local agencies have specific jurisdiction over paleontological resources and therefore, no state or local agencies were contacted.

#### **5.16.7 Permits Required and Permit Schedule**

No state or local agency requires a paleontological collecting permit to allow for the salvage of fossil remains discovered as a result of construction-related earth moving on non-federal public or private land in a project site.

#### **5.16.8 References**

- Adkison, W.L. 1973. Lithological characteristics of Upper Oligocene and Miocene rocks drilled at Elk Hills, Kern County, California. U.S. Geological Survey Bulletin 1375, 113 p.
- Anderson, F.M. 1905. A stratigraphic study in the Mount Diablo Range of California: California Academy of Science Proceedings, vol. 2, p. 155-248.
- Anderson, R., and R.W Pack. 1915. Geology and oil resources of the west border of the San Joaquin Valley, north of Coalinga, California: U.S. Geological Survey Bulletin 603, 220 p.
- Arnold, R., and H.R. Johnson. 1910. Preliminary report on the McKittrick-Sunset oil region, Kern and San Luis Obispo Counties, California: U.S. Geological Survey Bulletin 406.
- Bailey, E. H. (editor). 1966. Geology of northern California: California Division of Mines Bulletin 190, 508 p.
- Bartow, J.A. 1987. The Cenozoic evolution of the San Joaquin Valley, California: U.S. Geological Survey, Open-File Report OF-87-581, scale 1:500,000.

- Bartow, J.A. 1991. The Cenozoic evolution of the San Joaquin Valley, California: U.S. Geological Survey Professional Paper 1501, 40 p., scale 1:500,000.
- Bartow, J.A., and D.E. Marchand. 1979. Preliminary geologic map of Cenozoic deposits of the Clay area, California: U.S. Geological Survey Open-File Report 79-667 (scale 1:62,500).
- Berryman, W.M. 1973. Lithological characteristics of Pliocene rocks cored at Elk Hills, Kern County, California: Contributions to economic geology, U.S. Geological Survey Bulletin 1332-D, 56 p.
- Beyer, L.A., and A. Bartow. 1988. Summary of geology and petroleum plays to assess undiscovered recoverable petroleum resources, San Joaquin basin province, California: U.S. Geological Survey Open-File Report OF-87-450-Z, 80 p., scale 1:500,000.
- Bull, W.B. 1973. Geologic factors affecting compaction of deposits in a land-subsidence area [Fresno, Kings, and Merced counties, California]: Geological Society of America Bulletin, vol. 84, no. 12, p. 3783-3802.
- Bureau of Land Management (BLM). 1998. General procedural guidance for paleontological resource management: BLM Manual H-8270-1.
- California State Historic Preservation Office. 1983. Summary of state/federal laws protecting cultural resources: California State Historic Preservation Office, Sacramento, CA, 4 p.
- California Energy Commission (CEC). 2007. Rules of practice and procedure and power plant site certification regulations, April 2007: California Energy Commission, Sacramento, CA, 148 p.
- Callaway, D.C., and E.W. Rennie. 1991. San Joaquin basin, California: Geological Society of America, The Geology of North America, vol. P-2, Chapter 26, p. 417-430.
- Clark, B.L. 1929. Tectonics of the Valle Grande of California: American Association of Petroleum Geologists Bulletin, vol. 13, p. 199-238.
- Croft, M.G., and G.V. Gordon. 1968. Geology, hydrology, and quality of water in the Hanford – Visalia are, San Joaquin Valley, California: U.S. Geological Survey Open-File Report OF-68-67, 63 p., scale 1:125,000.
- Croft, M.G., and C. Wahrhaftig. 1965. General geology of the San Joaquin Valley: International Association of Quaternary Research, 7<sup>th</sup> Congress Guidebook, Field Conference I, Northern Great Basin and California, Nebraska Academy of Sciences, Lincoln, p. 133-137.
- Davis, G.H., and F.R. Hall. 1959. Water quality of eastern Stanislaus and northern Merced counties, California: Stanford University Publications, Geological Sciences, vol. 6, no. 1, p. 1-56.
- Davis, G.H., J.H. Green, F.H. Olmsted, and D.W. Brown. 1957. Groundwater conditions and storage capacity in the San Joaquin Valley, California: U.S. Geological Survey Open-File Report, 559 p.
- Davis, G.H., J.H. Green, F.H. Olmsted, and D.W. Brown. 1959. Groundwater conditions and storage capacity in the San Joaquin Valley, California: U.S. Geological Survey Water-Supply Paper 1469, 287 p.

- Davis, G.H., B.E. Lofgren, and S. Mack. 1964. Use of ground-water reservoirs for storage of surface water in the San Joaquin Valley, California: U.S. Geological Survey Water-Supply Paper 1618, 125 p.
- Dibblee, T.W. 1972. Geologic maps of fourteen 15-minute quadrangles along the San Andreas fault in the vicinity of Paso Robles and Cholame southeastward to Maricopa and Cuyama, California: U.S. Geological Survey Open-File Report OF-72-89, scale 1:62,500.
- Dibblee, T.W., Jr. 1973. Stratigraphy of the southern Coast Ranges near the San Andreas fault from Cholame to Maricopa, California: U.S. Geological Survey Professional Paper 764, 45 p.
- Dibblee, T.W. 2005a. Geologic map of the Taft and Mouth of Kern Quadrangles, Kern County, California: Dibblee Geological Foundation Map DF-95, scale 1:24,000.
- Dibblee, T.W. 2005b. Geologic map of the Fellows Quadrangle, Kern County, California: Dibblee Geological Foundation Map DF-96, scale 1:24,000.
- Dibblee, T.W. 2005c. Geologic map of the Panorama Hills Quadrangle, San Luis Obispo and Kern Counties, California: Dibblee Geological Foundation Map DF-97, scale 1:24,000.
- Dibblee, T.W. 2005d. Geologic map of the Reward Quadrangle, San Luis Obispo and Kern Counties, California: Dibblee Geological Foundation Map DF-100, scale 1:24,000.
- Dibblee, T.W. 2005e. Geologic map of the West Elk Hills Quadrangle, Kern County, California: Dibblee Geological Foundation Map DF-102, scale 1:24,000.
- Dibblee, T.W. 2005f. Geologic map of the East Elk Hills and Tupman Quadrangles, Kern County, California: Dibblee Geological Foundation Map DF-103, scale 1:24,000.
- Fenneman, N.M. 1931. Physiography of western United States: McGraw-Hill Book Company, New York, NY, 534 p.
- Fisk, L.H., and L.A. Spencer. 1994. Highway construction projects have legal mandates requiring protection of paleontologic resources (fossils), p. 213-225, *in* Burns, S.F. (editor), Proceedings of the 45<sup>th</sup> Highway Geology Symposium, Portland, OR, 258 p.
- Fisk, L.H., L.A. Spencer, and D.P. Whistler. 1994. Paleontologic resource impact mitigation on the PGT-PG&E Pipeline Expansion Project, Volume II: PG&E Section, California: unpublished report prepared for the Federal Energy Regulatory Commission, California Public Utilities Commission, Pacific Gas and Electric Company, and Bechtel Corporation by Paleo Environmental Associates, Inc., Altadena, CA, 123 p.
- Foss, C.D., and R. Blaisdell. 1968. Stratigraphy of the west side southern San Joaquin Valley, p. 33-43, *in* Karp, S. E. (editor), Guidebook: Geology and oil fields, west side southern San Joaquin Valley, 43<sup>rd</sup> annual meeting, Pacific Sections, American Association of Petroleum Geologists, Society of Exploration Geophysicists, and Society of Economic Paleontologists and Mineralogists.
- Gastaldo, R.A. 1999. International laws: collecting, transporting and ownership of fossils – USA, p. 330-338 *in* Jones, T.P. and N.P. Rowe (editors), Fossil plants and spores, The Geological Society, London, England, 396 p.

- Gester, G.C. 1917. Geology of a portion of the McKittrick district – a typical example of the west side San Joaquin Valley oil fields, and a correlation of the oil sands of the west side fields: California Academy of Sciences, 4<sup>th</sup> Series, vol. 7, no. 8, p. 207-227.
- Hackel, O. 1966. Summary of the geology of the Great Valley: p. 217-238 *in* E.H. Bailey (editor), Geology of Northern California: California Division of Mines and Geology Bulletin 190, 508 p.
- Harradine, F. 1950. Soils of western Fresno County, California: University of California, Agricultural Experiment Station, Division of Soils, Berkeley, CA, 86 p.
- Hoffman, R.D. 1964. Geology of the northern San Joaquin Valley: San Joaquin Geological Society Selected Papers, vol. 2, p. 30-45.
- Hoots, H.W., T.L. Bear, and W.D. Kleinpell. 1954. Geological summary of the San Joaquin Valley, California: p. 113-129 *in* Jahns, R.H. (editor), Geology of Southern California: California Division of Mines Bulletin 170, 289 p.
- Jahns, R.H. (editor). 1954. Geology of Southern California: California Division of Mines Bulletin 170, 289 p.
- Jefferson, G.T. 1991a. A catalogue of Late Quaternary vertebrates from California, part one, nonmarine lower vertebrate and avian taxa: Natural History Museum of Los Angeles County Technical Reports, no. 5, 60 p.
- Jefferson, G.T. 1991b. A catalogue of Late Quaternary vertebrates from California, Part Two, mammals: Natural History Museum of Los Angeles County Technical Reports, no. 7, 129 p.
- Jenkins, O.P. 1938. Geologic map of California: California Division of Mines and Geology, Sacramento, CA, 1:500,000 scale.
- Jennings, C.W. 1977. Geologic map of California: California Division of Mines and Geology, 1:750,000 scale.
- Lander, E.B. 1989. Interim paleontological resource technical report, Eastside Reservoir Project Study – Phase 1, Riverside County, California: unpublished report prepared for Metropolitan Water District of Southern California by Paleo Environmental Associates, Inc., Altadena, CA, 20 p.
- Lander, E.B. 1993. Paleontologic/cultural resource impact mitigation program final report: unpublished report prepared for Midway Sunset Cogeneration Company, Mojave Natural Gas Pipeline, and Kern County, California by Paleo Environmental Associates, Inc., Altadena, CA, 57 p.
- Lettis, W.R. 1982. Late Cenozoic stratigraphy and structure of the western margin of the central San Joaquin Valley, California: U.S. Geological Survey Open-File Report 82-526, 203 p.
- Lettis, W.R. 1988. Quaternary geology of the northern San Joaquin Valley: p. 333-351 *in* Graham, S.A., editor, Studies of the geology of the San Joaquin Basin, Pacific Section, Society of Economic Paleontologists and Mineralogists, vol. 60, 351 p.
- Lettis, W.R., and J.R. Unruh. 1991. Quaternary geology of the Great Valley, California: p. 164-176 *in* Dupré, W.R. and others, Quaternary geology of the Pacific margin, p. 141-213 *in*

- Morrison, R.B., editor, Quaternary nonglacial geology – conterminous U.S. Geological Society of America, Geology of North America, vol. K-2, 672 p.
- Maher, J.C., R.D. Carter, and R.J. Lantz. 1975. Petroleum geology of Naval Petroleum Reserve No. 1, Elk Hills, Kern County, California: U.S. Geological Survey Professional Paper 912.
- Maher, J.C., W.M. Trollman, and J.M. Denman. 1973. Geological literature on the San Joaquin Valley of California: Northern California Geological Society and Pacific Section of the American Association of Petroleum Geologists, Sacramento, CA, 582 p.
- Manning, J.C. 1968. Two late Pleistocene radiocarbon dates near Buttonwillow, California, p. 98-99 *in* Karp, S.E. (editor), Guidebook: Geology and oil fields, west side southern San Joaquin Valley. 43<sup>rd</sup> annual meeting, Pacific Sections, American Association of Petroleum Geologists, Society of Exploration Geophysicists, and Society of Economic Paleontologists and Mineralogists, 142 p.
- Marchand, D.E. 1977. The Cenozoic history of the San Joaquin Valley and the adjacent Sierra Nevada as inferred from the geology and soils of the eastern San Joaquin Valley: p. 39-50 *in* Singer, M.J. (editor), Soil development, geomorphology, and Cenozoic history of the northeastern San Joaquin Valley and adjacent areas, California: University of California Press, Guidebook for Joint Field Session, Soil Science Society of America and Geological Society of America, 328 p.
- Marchand, D.E., and A. Allwardt. 1981. Late Cenozoic stratigraphic units, northeastern San Joaquin Valley, California: U.S. Geological Survey Bulletin 1470, 70 p.
- Marshall, L.G. 1976. Paleontological salvage and federal legislation: Journal of Paleontology, vol. 50, p. 346-348.
- Mendenhall, W.C. 1908. Preliminary report on the ground waters of San Joaquin Valley, California: U.S. Geological Survey Water-Supply Paper 222, 52 p.
- Mendenhall, W.C., R.B. Dole, and H. Stabler. 1916. Ground water in San Joaquin Valley, California: U.S. Geological Survey Water-Supply Paper, 310 p.
- Merriam, J.C. 1903. The Pliocene and Quaternary Canidae of the Great Valley of California: University of California Publications, Bulletin of the Department of Geology, vol. 3, no. 14, p. 277-290.
- Merriam, J.C. 1905. A new sabre-tooth from California: University of California Publications, Bulletin of the Department of Geology, vol. 4, no. 9, p. 171-175.
- Merriam, J.C. 1914. The Tertiary of the Great Basin and that of the marginal marine province in California: Science, vol. 40, p. 643-645.
- Merriam, J.C. 1915a. Tertiary vertebrate faunas of the North Coalinga region of California – a contribution to the study of paleontologic correlation in the Great Basin and Pacific Coast Provinces: Transactions of the American Philosophical Society, vol. 22, no. 3, p. 191-234.
- Merriam, J.C. 1915b. Faunas of the San Pablo Bay syncline and the Mount Diablo region: Meeting of the Paleontological Society Field Guide, 2 - 7 August 1915, 15 p.

- Merriam, J.C. 1917. Relationships of Pliocene mammalian faunas from the Pacific Coast and Great Basin Provinces of North America: University of California Publications, Bulletin of the Department of Geology, vol. 10, no. 22, p. 421-443.
- Miller, R.E., J.H. Green, and G.H. Davis. 1971. Geology of the compacting deposits in the Los Banos – Kettleman City subsidence area, California -- mechanics of aquifer systems: U.S. Geological Survey Professional Paper 497-E. 45 p.
- Morton, P.K., and B.W. Troxel. 1962. Mines and mineral resources of Kern County, California: California Division of Mines and Geology County Report 1, 370 p, scale 1:320,000.
- Page, R.W. 1986. Geology of the fresh ground-water basin of the Central Valley, California, with texture maps and sections: U.S. Geological Survey Professional Paper 1401-C, 54 p., scale 1:500,000.
- Piper, A.M., H.S. Gale, H.E. Thomas, and T.W. Robinson. 1939. Geology and ground-water hydrology of the Mokelumne area, California: U.S. Geological Survey Water-Supply Paper 780, 230 p.
- Porter, L.E. 1943. Elk Hills Oil and Gas Field (U.S. Naval Petroleum Reserve No. 1), p. 512-516, *in* Jenkins, O.P. (compiler), Geologic formations and economic development of the oil and gas fields of California: California Division of Mines Bulletin 118, 516 p.
- Repenning, C.A. 1980. Paleontologic resource evaluation, Naval Petroleum Reserve No. 1 (Elk Hills): U.S. Geological Survey Branch of Paleontology and Stratigraphy, Menlo Park, California.
- Reynolds, R.E. 1987. Paleontologic resource assessment, Midway-Sunset Cogeneration Project, Kern County, California: unpublished report prepared for Southern California Edison Company by San Bernardino County Museum, San Bernardino, CA, 15 p.
- Reynolds, R.E. 1990. Paleontological mitigation program, Midway-Sunset Cogeneration Project, Kern County, California: unpublished report prepared for Midway-Sunset Cogeneration Company, by San Bernardino County Museum, San Bernardino, CA, 45 p.
- Shipman, P. 1977. Paleoecology, taphonomic history and population dynamics of the vertebrate assemblage from the middle Miocene of Fort Turnan, Kenya: unpublished PhD dissertation, New York University, New York, 193 p.
- Shipman, P. 1981. Spatial distribution of fossils in sediments: p. 65-98 *in* Shipman, P., Life history of a fossil, an introduction to taphonomy and paleoecology, Harvard University Press, Cambridge, MA, 222 p.
- Smith, A.R. 1964. Geologic map of California – Bakersfield sheet: California Division of Mines and Geology, scale 1:250,000.
- Society of Vertebrate Paleontology. 1995. Assessment and mitigation of adverse impacts to nonrenewable paleontologic resources – standard guidelines: Society of Vertebrate Paleontology News Bulletin, vol. 163, p. 22-27.
- Society of Vertebrate Paleontology. 1996. Conditions of receivership for paleontologic salvage collections: Society of Vertebrate Paleontology News Bulletin, vol. 166, p. 31-32.

- Spencer, L.A. 1990. Paleontological mitigation program, Midway-Sunset Cogeneration Project, Natural Gas Pipeline, Kern County, California: unpublished report prepared for Midway-Sunset Cogeneration Company by Paleo Environmental Associates, Inc., Altadena, CA, 12 p.
- Stirton, R.A., and V.L. VanderHoof. 1933. Osteoborus, a new genus of dogs, and its relations to Borophagus Cope: University of California Publications, Bulletin of the Department of Geological Sciences, vol. 23, no. 4, p. 175-182.
- Taylor, D.W. 1966. Summary of North American Blancan nonmarine mollusks: Malacologia vol. 4, no. 1, 172 p.
- Wahrhaftig, C., S.W. Stine, and N.K. Huber. 1993. Quaternary geologic map of the San Francisco Bay 4° x 6° Quadrangle, United States: U.S. Geological Survey Miscellaneous Investigations Map I-1420, 1:1,000,000 scale.
- Wells, J.C. 1952. Elk Hills Field, California, p. 241-245, *in* Guidebook Field Trip Routes, Geology, Oil Fields: American Association of Petroleum Geologists, Society of Exploration Geophysicists, and Society of Economic Paleontologists and Mineralogists, Pacific Section Guidebook Joint Annual Meeting, Los Angeles, California, March 1952, 290 p.
- West, R.M. 1991. State regulation of geological, paleontological, and archaeological collecting: Curator, vol. 34, p. 199-209.
- White, R.E. 1987. Paleomagnetism of the Tulare Formation: unpublished masters thesis, California State University, Longbeach, CA. 272 p.
- Wood, P.R., and G.H. Davis. 1959. Ground-water conditions in the Avenal-McKittrick area, Kings and Kern Counties, California: U.S. Geological Survey Water-Supply Paper 1457.
- Woodring, W.P., P.V. Roundy, and H.E. Farnsworth. 1932. Geology and oil resources of the Elk Hills, California, including Naval Petroleum Reserve No. 1: U.S. Geological Survey Bulletin 835.
- Woodring, W.P., R. Stewart, and R.W Richards. 1940. Geology of the Kettleman Hills oil field, California; U.S. Geological Survey Professional Paper 195.

*This page intentionally left blank*

Adequacy Issue: Adequate Inadequate DATA ADEQUACY WORKSHEET Revision No. 0 Date \_\_\_\_\_

Technical Area: **Paleontological Resources** Project: \_\_\_\_\_ Technical Staff: \_\_\_\_\_

Project Manager: \_\_\_\_\_ Docket: \_\_\_\_\_ Technical Senior: \_\_\_\_\_

| SITING REGULATIONS      | INFORMATION  | AFC PAGE NUMBER AND SECTION NUMBER            | ADEQUATE YES OR NO | INFORMATION REQUIRED TO MAKE AFC CONFORM WITH REGULATIONS |
|-------------------------|--|---|--------------------|---|
| Appendix B (g) (1)      | ...provide a discussion of the existing site conditions, the expected direct, indirect and cumulative impacts due to the construction, operation and maintenance of the project, the measures proposed to mitigate adverse environmental impacts of the project, the effectiveness of the proposed measures, and any monitoring plans proposed to verify the effectiveness of the mitigation.                              | Sections 5.16.1.1, 5.16.2, 5.16.3, and 5.16.4 |                    |   |
| Appendix B (g) (16) (A) | Identification of the physiographic province and a brief summary of the geologic setting, formations, and stratigraphy of the project area. The size of the paleontological study area may vary depending on the depositional history of the region.   | Sections 5.16.1.1, 5.16.1.2, and 5.16.1.5.    |                    |   |
| Appendix B (g) (16) (B) | A discussion of the sensitivity of the project area described in subsection (g)(16)(A) and the presence and significance of any known paleontologic localities or other paleontologic resources within or adjacent to the project. Include a discussion of sensitivity for each geologic unit identified on the most recent geologic map at a scale of 1:24,000. Provide rationale as to why the sensitivity was assigned. | Section 5.16.1.5                              |                    |   |

Adequacy Issue: Adequate Inadequate DATA ADEQUACY WORKSHEET Revision No. 0 Date \_\_\_\_\_

Technical Area: **Paleontological Resources** Project: \_\_\_\_\_ Technical Staff: \_\_\_\_\_

Project Docket: \_\_\_\_\_ Technical Senior: \_\_\_\_\_

Manager: \_\_\_\_\_

| SITING REGULATIONS      | INFORMATION  | AFC PAGE NUMBER AND SECTION NUMBER           | ADEQUATE YES OR NO | INFORMATION REQUIRED TO MAKE AFC CONFORM WITH REGULATIONS |
|-------------------------|--|--|--------------------|---|
| Appendix B (g) (16) (C) | A summary of all local museums, literature searches and field surveys used to provide information about paleontologic resources in the project area described in subsection (g)(16)(A). Identify the dates of the surveys, methods used in completing the surveys, and the names and qualifications of the individuals conducting the surveys.   | Section 5.16.1.3                             |                    |   |
| Appendix B (g) (16) (D) | Information on the specific location of known paleontologic resources, survey reports, locality records, and maps at a scale of 1:24,000, showing occurrences of fossil finds, if known, within a one-mile radius of the project and related facilities shall be included in a separate appendix to the Application and submitted to the Commission under a request for confidentiality, pursuant to Title 20, California Code of Regulations, s 2501 <i>et seq.</i> | Appendix Q, Paleontological Technical Report |                    |   |
| Appendix B (g) (16) (E) | A discussion of any educational programs proposed to enhance awareness of potential impacts to paleontological resources by employees, measures proposed for mitigation of impacts to known paleontologic resources, and a set of contingency measures for mitigation of potential impacts to currently unknown paleontologic resources.   | Section 5.16.4                               |                    |   |

Adequacy Issue: Adequate Inadequate DATA ADEQUACY WORKSHEET Revision No. 0 Date \_\_\_\_\_

Technical Area: **Paleontological Resources** Project: \_\_\_\_\_ Technical Staff: \_\_\_\_\_

Project Manager: \_\_\_\_\_ Docket: \_\_\_\_\_ Technical Senior: \_\_\_\_\_

| SITING REGULATIONS        | INFORMATION  | AFC PAGE NUMBER AND SECTION NUMBER | ADEQUATE YES OR NO | INFORMATION REQUIRED TO MAKE AFC CONFORM WITH REGULATIONS |
|---------------------------|--|------------------------------------|--------------------|---|
| Appendix B<br>(i) (1) (A) | Tables which identify laws, regulations, ordinances, standards, adopted local, regional, state, and federal land use plans, leases, and permits applicable to the proposed project, and a discussion of the applicability of, and conformance with each. The table or matrix shall explicitly reference pages in the application wherein conformance, with each law or standard during both construction and operation of the facility is discussed; and | Table 5.16-1; Section 5.16.5       |                    |   |
| Appendix B<br>(i) (1) (B) | Tables which identify each agency with jurisdiction to issue applicable permits, leases, and approvals or to enforce identified laws, regulations, standards, and adopted local, regional, state and federal land use plans, and agencies which would have permit approval or enforcement authority, but for the exclusive authority of the commission to certify sites and related facilities.  | Table 5.16-1; Section 5.16.5       |                    |   |
| Appendix B<br>(i) (2)     | The name, title, phone number, address (required), and email address (if known), of an official who was contacted within each agency, and also provide the name of the official who will serve as a contact person for Commission staff.   | N/A                                |                    |   |
| Appendix B<br>(i) (3)     | A schedule indicating when permits outside the authority of the commission will be obtained and the steps the applicant has taken or plans to take to obtain such permits.   | Section 5.16.7                     |                    |   |

